

Growth of fat slits and dispersionless KP hierarchy

A. Zabrodin *

November 2008

Abstract

A “fat slit” is a compact domain in the upper half plane bounded by a curve with endpoints on the real axis and a segment of the real axis between them. We consider conformal maps of the upper half plane to the exterior of a fat slit parameterized by harmonic moments of the latter and show that they obey an infinite set of Lax equations for the dispersionless KP hierarchy. Deformation of a fat slit under changing a particular harmonic moment can be treated as a growth process similar to the Laplacian growth of domains in the whole plane. This construction extends the well known link between solutions to the dispersionless KP hierarchy and conformal maps of slit domains in the upper half plane and provides a new, large family of solutions.

*Institute of Biochemical Physics, 4 Kosygina st., 119334, Moscow, Russia and ITEP, 25 B.Chermushkinskaya, 117218, Moscow, Russia

1 Introduction

Parametric families of conformal maps in 2D are known to be closely related to long wave limits of nonlinear integrable PDE's and their infinite hierarchies. This observation was first made in [1] for mappings of slit domains and then extended to mappings of domains bounded by Jordan curves in [2]. In both cases conformal maps from a standard reference domain (such as upper half plane, or unit disk) to a domain of a varying shape serve as Lax functions of an integrable hierarchy whose flows are identified with variations of the conformal maps described by an infinite set of Lax equations. The integrable structures arising in this way are dispersionless Kadomtsev-Petviashvili (dKP) and dispersionless 2D Toda (dToda) hierarchies and, more generally, the universal Whitham hierarchy first introduced [3, 4] in an absolutely different context with the aim to describe slow modulations of exact solutions to soliton equations.

The most promising progress along these lines was achieved in geometric and physical interpretation of the dToda hierarchy. The key fact, established in [5] and further elaborated in [6, 7], is that variations of domains under the Toda flows go exactly according to the Darcy law specific for growth processes of Laplacian type and viscous hydrodynamics in the Hele-Shaw cell with zero surface tension (see, e.g., [8, 9]).

Although the dKP hierarchy is simpler than the dToda hierarchy, its role in the theory of conformal maps and Laplacian growth is not well understood. The connection with conformal maps observed in [1] gives a geometric interpretation to only rather special (degenerate) solutions of the dKP hierarchy which are reductions to systems of hydrodynamic type with a finite number of degrees of freedom. As is shown in [1] (see also [10, 11]), they are related to conformal maps of the upper half plane with slits emanating from the real axis.

The aim of this paper is to demonstrate that the geometric interpretation of the dKP hierarchy is not limited by domains of such special kind. We show that the same dKP hierarchy is able to cover a much broader class of domains which can be obtained from the upper half plane by removing not just an infinitely thin slit but a whole compact piece (of a non-zero area and arbitrary shape) attached to the real axis, which we call a “fat slit” to stress the analogy (Fig. 1). There is an important difference, however. The dKP-evolution of usual slits is actually finite dimensional because the slits are to be regarded as arcs of fixed curves [1, 10], so only their endpoints can move. In contrast, the dKP-evolution of fat slits (given by the Lax equations) takes place in an infinite dimensional variety corresponding to changing their shape in an arbitrary way.

Let us recall the Lax formulation of the dKP hierarchy. Starting from a Laurent series

$$z(p) = p + \sum_{k=1}^{\infty} a_k p^{-k} \quad (1.1)$$

one introduces the dependence on an infinite number of “times” T_1, T_2, T_3, \dots via Lax equations

$$\frac{\partial z(p)}{\partial T_k} = \{B_k(p), z(p)\} := \frac{\partial B_k}{\partial p} \frac{\partial z}{\partial T_1} - \frac{\partial B_k}{\partial T_1} \frac{\partial z}{\partial p}, \quad (1.2)$$

where the generators of the flows B_k are polynomials in p of the form $B_k(p) = (z^k(p))_{\geq 0}$ (polynomial parts of $z^k(p)$). The dKP hierarchy is an infinite system of nonlinear PDE's




Figure 1: a) A slit; b) A “fat slit”.

for a_i ’s resulting from comparing coefficients in front of different powers of p in the Lax equations or in an equivalent system of equations of the Zakharov-Shabat type

$$\frac{\partial B_j(p)}{\partial T_k} - \frac{\partial B_k(p)}{\partial T_j} + \{B_j(p), B_k(p)\} = 0 \quad (1.3)$$

for all $j, k \geq 1$ (the Poisson bracket is defined in (1.2)). The coefficients a_i and the times T_i are assumed to be real numbers.

Assuming that $z(p)$ is a normalized conformal map from the upper half plane onto the exterior of a fat slit, we show that it obeys equations (1.2), where T_k are properly defined harmonic moments of the fat slit (or rather of its exterior). This fact follows from the Hadamard formula for variations of the Green function with Dirichlet boundary conditions. In this sense the arguments are parallel to [12, 13]. Evolution in $T = T_1$ with all other times fixed has an interpretation as a version of the Laplacian growth in the upper half plane with fixed real axis.

Applying the approach developed in [12, 13, 14] to the case of fat slits in the upper half plane, we construct the dispersionless “tau-function” (which is actually a limit of properly rescaled logarithm of a dispersionfull tau-function) of the dKP hierarchy as a functional on the space of fat slits explicitly given by

$$\log \tau = -\frac{1}{\pi^2} \iint_{\text{fat slit}} \log \left| \frac{z - \zeta}{z - \bar{\zeta}} \right| d^2 z d^2 \zeta. \quad (1.4)$$

It has a clear electrostatic interpretation as Coulomb energy of a fat slit filled with electric charge of a uniform density in the presence of an infinite grounded conductor placed along the real axis. This functional regarded as a function of harmonic moments obeys a dispersionless version of the Hirota relation which serves as a master equation generating the whole dKP hierarchy.

2 Fat slit domains, their conformal maps and Green’s functions

Consider a compact simply connected domain \mathbf{B} in the upper half-plane \mathbb{H} bounded by a non-self-intersecting analytic curve γ in \mathbb{H} with endpoints x_- , x_+ on the real axis and a segment of the real axis between them. This segment will be called the *base* of \mathbf{B} . Without loss of generality, one can assume that the origin belongs to the base. For brevity, and in




Figure 2: A fat slit B in the upper half plane and the corresponding symmetric domain $D = B \cup \bar{B}$.

order to emphasize an analogy with slit domains manifested in the common integrable structure of their conformal maps, we call such a domain a *fat slit*. Accordingly, the complement, $\mathbb{H} \setminus B$, will be referred to as a domain with a fat slit, or simply a fat slit domain (in our case, the fat slit half-plane).

It is often convenient to treat fat slits as upper halves of domains symmetric with respect to the real axis. Namely, set $D = B \cup \bar{B}$, where \bar{B} is the domain in the lower half plane which is obtained from B by complex conjugation $z \rightarrow \bar{z}$ (Fig. 2). Obviously, the domain D is symmetric with respect to the complex conjugation. In what follows we call such domains simply *symmetric*. Vice versa, any compact simply connected symmetric domain D is a union of a fat slit and its complex conjugate. The boundary of D is assumed to be analytic everywhere except the two points on the real axis which are allowed to be corner points.

2.1 Conformal maps

Let $p(z)$ be a conformal map from $\mathbb{H} \setminus B$ (in the z -plane) onto \mathbb{H} (in the p -plane) shown schematically in Fig. 3. We normalize it by the condition that the expansion of $p(z)$ in a Laurent series at infinity is of the form

$$p(z) = z + \frac{u}{z} + \sum_{k \geq 2} u_k z^{-k}, \quad |z| \rightarrow \infty \quad (2.1)$$

(a “hydrodynamic” normalization). Assuming this normalization, the map is unique. The upper part of the boundary, γ , is mapped to a segment of the real axis $[p_-, p_+]$, while the rays of the real axis outside B are mapped to the real rays $[-\infty, p_-]$ and $[p_+, \infty]$ (Fig. 3). From this it follows that the coefficients u_k are all real numbers. The first coefficient, $u_1 := u$, is called a *capacity* of B . It is known to be positive. We also




Figure 3: *The conformal map $p(z)$.*

need the inverse map, $z(p)$, which can be expanded into the inverse Laurent series

$$z(p) = p - \frac{u}{p} + \sum_{k=2}^{\infty} a_k p^{-k}, \quad |p| \rightarrow \infty \quad (2.2)$$

with real coefficients a_k connected with u_k by polynomial relations. The series converges for large enough $|p|$.

According to the Schwarz symmetry principle, the function $z(p)$ admits an analytic continuation to the lower half plane. This analytically continued function performs a conformal map from the whole complex plane with a finite cut on the real axis from p_- to p_+ onto the exterior of the symmetric domain $D = B \cup \bar{B}$.

2.2 Green's functions

Let $D = B \cup \bar{B}$ be a symmetric domain and let $G(z, z')$ be the standard Green's function of the Dirichlet boundary problem in $\mathbb{C} \setminus D$. The function is harmonic in $\mathbb{C} \setminus D$ with respect to both variables except at $z = z'$, where it has a logarithmic singularity $G(z, z') = \log |z - z'| + \dots$ and equals zero when either z or z' lies on the boundary of D . The Green's function solves the Dirichlet boundary value problem: the formula

$$f(z) = -\frac{1}{2\pi} \oint_{\partial D} f(\xi) \partial_n G(z, \xi) |d\xi| \quad (2.3)$$

harmonically extends the function $f(\xi)$ from the contour ∂D to its exterior. Here and below, ∂_n is the normal derivative at the boundary, with the normal vector being directed to the exterior of D . We also note the Hadamard formula [15] for variation of the Green's function under variation of the domain:

$$\delta G(a, b) = \frac{1}{2\pi} \oint_{\partial D} \partial_n G(a, z) \partial_n G(b, z) \delta n(z) |dz|, \quad (2.4)$$

where $\delta n(z)$ is the infinitesimal normal displacement of the contour. Some care is needed to define a deformation near the corner points. However, for our purposes it is enough to consider deformations with fixed corner points, then $\delta n(z)$ is well defined at any point of the boundary. Also, in this paper we consider only the case when both angles α_+ and α_- are acute, $0 < \alpha_{\pm} < \pi/2$, then the normal derivative of the Green's function vanishes at the corners and the integral converges (see more details below).

For symmetric domains the Green's function obeys the property $G(z, z') = G(\bar{z}, \bar{z}')$. Let us call a function $f(z)$ *even* (respectively, *odd*) if $f(z) = f(\bar{z})$ (respectively, $f(z) = -f(\bar{z})$).

$-f(\bar{z})$). It is natural to introduce even and odd Green's functions such that $G^\pm(z, z') = \pm G^\pm(z, \bar{z}')$:

$$G^\pm(z, z') = G(z, z') \pm G(z, \bar{z}').$$

Note that $G^-(x, z) = 0$ for real x . The Poisson formula (2.3) for even and odd boundary functions can be written in the form

$$f(z) = -\frac{1}{2\pi} \int_\gamma f(\xi) \partial_n G^\pm(z, \xi) |d\xi|, \quad (2.5)$$

where the integration goes over the *non-closed* contour γ (Fig. 4). The Hadamard formula for G^\pm ,

$$\delta G^\pm(a, b) = \frac{1}{2\pi} \int_\gamma \partial_n G^\pm(a, z) \partial_n G^\pm(b, z) \delta n(z) |dz| \quad (2.6)$$

directly follows from (2.4), taking into account that deformations of symmetric domains obey the condition $\delta n(z) = \delta n(\bar{z})$.

2.3 The odd Green's function

An important part in what follows is played by the odd Green's function G^- . It solves the following Dirichlet boundary value problem in \mathbb{H} : To find a harmonic function in $\mathbb{H} \setminus \mathbf{B}$ bounded at infinity such that it is equal to a given function on γ and 0 on the rays of the real axis outside \mathbf{B} . Similar to the Green's function G , G^- can be expressed through a conformal map to a fixed reference domain. The most natural reference domain in our case is the upper half plane \mathbb{H} . It is easy to see that

$$G^-(z, z') = \log \left| \frac{p(z) - p(z')}{p(z) - p(\bar{z}')} \right|, \quad (2.7)$$

where $p(z)$ is the conformal map (2.1) from $\mathbb{H} \setminus \mathbf{B}$ onto \mathbb{H} . We also need a useful formula for the kernel $\partial_n G^-(a, z)$ in (2.5) through the conformal map,

$$\partial_n G^-(a, z) = -2 \operatorname{Im} p(a) \frac{|p'(z)|}{|p(z) - p(a)|^2}, \quad z \in \gamma \quad (2.8)$$

(which straightforwardly follows from (2.7)) and its limiting case as $|a| \rightarrow \infty$:

$$\partial_n \operatorname{Im} p(z) = |p'(z)|, \quad z \in \gamma. \quad (2.9)$$

Let us present the expansion of the odd Green's function $G^-(a, z)$ as $|a| \rightarrow \infty$:

$$G^-(a, z) = 2 \sum_{k \geq 1} \frac{1}{k} \operatorname{Im}(a^{-k}) \operatorname{Im}(B_k(p(z))). \quad (2.10)$$

Here $B_k(p)$ are Faber polynomials of $p(z)$ defined by the expansion

$$\log \frac{z}{p(z) - p} = \sum_{k \geq 1} \frac{z^{-k}}{k} B_k(p), \quad |z| \rightarrow \infty, \quad (2.11)$$

and explicitly given by

$$B_k(p) = (z^k(p))_{\geq 0}, \quad (2.12)$$

where $(\dots)_{\geq 0}$ means the polynomial part of the Laurent series. Indeed, fixing a point $z_1 \in \mathbb{H} \setminus \mathbf{B}$, we have

$$\sum_{k \geq 1} \frac{z_1^{-k}}{k} B_k(p) = \sum_{k \geq 1} \frac{(z^k(p))_{\geq 0}}{k z^k(p_1)} = - \left[\log \left(1 - \frac{z(p)}{z(p_1)} \right) \right]_{\geq 0}$$

where $p_1 = p(z_1)$. To separate non-negative part, we write

$$\log \left(1 - \frac{z(p)}{z(p_1)} \right) = \log \frac{p_1 - p}{z(p_1)} + \log \frac{z(p_1) - z(p)}{p_1 - p}$$

and notice that the expansion of the first (second) term contains only non-negative (respectively, negative) powers of p . Therefore,

$$\sum_{k \geq 1} \frac{z_1^{-k}}{k} B_k(p) = -\log(p(z_1) - p) + \log z_1,$$

which coincides with (2.11). In particular, $B_1(p) = p$. Clearly,

$$B_k(p(z)) = z^k + O(z^{-1}), \quad |z| \rightarrow \infty, \quad (2.13)$$

and the function $B_k(p(z)) - z^k$ is analytic in $\mathbb{C} \setminus \mathbf{D}$.

3 Local coordinates in the space of fat slits

We are going to show that the harmonic moments T_k (defined as in (3.1) below) locally characterize a fat slit in the following sense. First, any small deformation of \mathbf{B} that preserves the moments T_k , is trivial, i.e., any non-trivial deformation changes at least one of them. This fact means local uniqueness of a fat slit having given moments. Second, the moments T_k , under certain conditions discussed below, are independent quantities meaning that one can explicitly define infinitesimal deformations of \mathbf{B} that change any one of them keeping all other fixed. In this weak sense they serve as local coordinates in the space of fat slits (c.f. the remark in Section 2.1 in [14]).

3.1 Harmonic moments

Given a fat slit \mathbf{B} , let us introduce harmonic moments of the fat slit domain $\mathbb{H} \setminus \mathbf{B}$ as

$$\begin{aligned} T_k &= \frac{2}{\pi k} \operatorname{Im} \int_{\mathbb{H} \setminus \mathbf{B}} z^{-k} d^2 z, \quad k \geq 2, \\ T_1 &= -\frac{2}{\pi} \operatorname{Im} \int_{\mathbf{B}} z^{-1} d^2 z. \end{aligned} \quad (3.1)$$

Here we assume that the base of \mathbf{B} is a segment containing zero. Since $\operatorname{Im}(z^{-1}) < 0$ for $z \in \mathbb{H}$, T_1 is always positive. Although the integrand in the formula for T_1 is singular at the origin, the integral converges. The integral for T_2 diverges at infinity, so one should




Figure 4: *The integration contour γ .*

introduce a cut-off at some large radius and make the angular integration first; this prescription is equivalent to the contour integral representation given below. Note that this set of moments does not include the area of \mathbf{B} . Note also that the standard harmonic moments dealt with in [13] are, for symmetric domains, real parts of the integrals in (3.1) rather than imaginary ones.

Some other integral representations of the moments (3.1) are also useful. Imaginary parts of the integrals can be taken by extending the integration to the lower half plane as

$$T_k = \frac{1}{i\pi k} \int_{\mathbb{C} \setminus \mathbf{D}} \text{sign}(y) z^{-k} d^2 z, \quad k \geq 2, \quad (3.2)$$

$$T_1 = -\frac{1}{i\pi} \int_{\mathbf{D}} \text{sign}(y) z^{-1} d^2 z,$$

where $y = \mathcal{I}m z$. Contour integral representations are easily obtained using the Stokes theorem. They read

$$T_k = \frac{2}{\pi k} \mathcal{I}m \int_{\gamma} y z^{-k} dz = \frac{1}{\pi i k} \oint_{\partial \mathbf{D}} |y| z^{-k} dz, \quad k \geq 1. \quad (3.3)$$

The non-closed integration contour γ (shown in Fig. 4) is the part of the boundary of \mathbf{B} lying in the upper half plane (with the orientation from right to left).

It is convenient to introduce the generating function of the moments T_k :

$$M_+(z) = \frac{1}{\pi i} \oint_{\partial \mathbf{D}} \frac{|y'| dz'}{z' - z} = \sum_{k=1}^{\infty} k T_k z^{k-1}, \quad |z| \rightarrow 0 \quad (3.4)$$

(here $y' = \mathcal{I}m z'$). The integral of Cauchy type in (3.4) defines an analytic function everywhere inside \mathbf{D} . In a small enough neighborhood of the origin this function is represented by the (convergent) Taylor series standing in the r.h.s. of (3.4).

For example, let \mathbf{B} be the half-disk of radius R : $|z| \leq R$, $\mathcal{I}m z \geq 0$. Then an easy calculation gives

$$T_k = \frac{4R^{2-k}}{\pi k^2(2-k)} \quad \text{for odd } k, \quad T_k = 0 \quad \text{for even } k$$

and

$$M_+(z) = \frac{R}{\pi} \left(2 + \frac{z^2 - R^2}{Rz} \log \frac{R - z}{R + z} \right).$$

3.2 An electrostatic interpretation

Similar to the standard harmonic moments from papers [2, 13], the moments T_k have a clear 2D electrostatic interpretation. Let the interior of the domain \mathbf{B} be filled by an electric charge with uniform density -1 and let the lower half plane (or just the real axis) be a grounded conductor. By the reflection principle, the electric potential $\Phi^-(z)$ in the upper half plane is equal to the potential created by the charge in \mathbf{B} and the fictitious “mirror” charge of opposite sign in $\bar{\mathbf{B}}$:

$$\Phi^-(z) = -\frac{2}{\pi} \int_{\mathbf{B}} \log \left| \frac{z - z'}{z - \bar{z}'} \right| d^2 z'. \quad (3.5)$$

Let us show that the T_k 's are coefficients in the multipole expansion of $\Phi^-(z)$ in the interior of \mathbf{B} near the origin. We have, for $z \in \mathbf{B}$:

$$\begin{aligned} \partial_z \Phi^-(z) &= \frac{1}{\pi} \int_{\mathbf{B}} \frac{d^2 z'}{z' - z} - \frac{1}{\pi} \int_{\bar{\mathbf{B}}} \frac{d^2 z'}{z' - z} \\ &= -\bar{z} + \frac{1}{2\pi i} \oint_{\partial \mathbf{B}} \frac{\bar{z}' dz'}{z' - z} - \frac{1}{2\pi i} \oint_{\partial \bar{\mathbf{B}}} \frac{\bar{z}' dz'}{z' - z} \\ &= z - \bar{z} - \frac{1}{\pi} \oint_{\partial \mathbf{B}} \frac{y' dz'}{z' - z} + \frac{1}{\pi} \oint_{\partial \bar{\mathbf{B}}} \frac{y' dz'}{z' - z}, \end{aligned}$$

where we substituted $\bar{z}' = z' - 2iy'$ in the second line and integrated the analytic parts by taking residues. Since the function under the integrals vanishes on the parts of the contours along the real axis, we can eliminate them and combine the two integrals into a single integral over $\partial(\mathbf{B} \cup \bar{\mathbf{B}}) = \partial \mathbf{D}$:

$$\partial_z \Phi^-(z) = z - \bar{z} - \frac{1}{\pi} \oint_{\partial \mathbf{D}} \frac{|y'| dz'}{z' - z} = z - \bar{z} - i \sum_{k \geq 1} k T_k z^{k-1}, \quad (3.6)$$

where the second equality follows from (3.4). Since $\Phi^-(x) = 0$ at real x , we obtain the expansion of the $\Phi^-(z)$ around 0 in the upper half plane:

$$\Phi^-(z) = \frac{1}{2}(z - \bar{z})^2 - i \sum_{k \geq 1} T_k (z^k - \bar{z}^k). \quad (3.7)$$

Similarly, expanding $\Phi^-(z)$ around ∞ in the upper half plane, we get:

$$\partial_z \Phi^-(z) = -i \sum_{k \geq 1} V_k z^{-k-1}, \quad \Phi^-(z) = i \sum_{k \geq 1} \frac{V_k}{k} (z^{-k} - \bar{z}^{-k}), \quad (3.8)$$

where

$$V_k = \frac{2}{\pi} \mathcal{I}m \int_{\mathbf{B}} z^k d^2 z \quad (3.9)$$

are moments of the interior. Their generating function, $M_-(z)$, is given by the same Cauchy-type integral (3.4) for z outside \mathbf{D} :

$$M_-(z) = \frac{1}{\pi i} \oint_{\partial \mathbf{D}} \frac{|y'| dz'}{z' - z} = \sum_{k=1}^{\infty} V_k z^{-k-1}, \quad |z| \rightarrow \infty. \quad (3.10)$$

Sometimes an equivalent electrostatic interpretation appears to be more convenient. Let us assume that there is no conductor but the domain $\bar{\mathbb{B}}$ in the lower half plane is indeed filled with the “mirror” charge. In this case formulas (3.6), (3.7) and (3.8) admit continuation to the lower half plane which is achieved by complex conjugation of both sides. For example, complex conjugation of (3.6) yields $\partial_{\bar{z}}\Phi^-(\bar{z}) = \bar{z} - z + i \sum_{k \geq 1} k T_k \bar{z}^{k-1}$, $z \in \mathbb{B}$, which can be rewritten as $\partial_z\Phi^-(z) = z - \bar{z} + i \sum_{k \geq 1} k T_k z^{k-1}$, $z \in \bar{\mathbb{B}}$.

3.3 Local uniqueness of a fat slit with given moments

Here, we prove the local uniqueness of a fat slit with given moments. The deformations changing only one moment will be constructed in the next subsection.

For the purpose of this section it is convenient to work with symmetric domains $\mathbb{D} = \mathbb{B} \cup \bar{\mathbb{B}}$ rather than with fat slits themselves. Let $\mathbb{D}(t)$ be a one-parameter deformation in the class of symmetric domains such that $\mathbb{D}(0) = \mathbb{D}$ and $\partial_t T_k = 0$ for all $k = 1, 2, \dots$. We shall show that any such deformation is trivial: $\mathbb{D}(t) = \mathbb{D}(0)$ (at least in a small neighborhood of $t = 0$). To see this, consider the t -derivative of the function $M_+(z)$. A simple calculation (see Appendix A) shows that

$$\partial_t M_+(z) = \frac{1}{\pi i} \oint_{\partial \mathbb{D}} \frac{\text{sign}(y') v_n(z')}{z - z'} |dz'|, \quad (3.11)$$

where $v_n(z) = \delta n(z)/\delta t$ is the “velocity” of the normal displacement of the boundary at the point z . (If $x(\sigma, t)$, $y(\sigma, t)$ is any parametrization of the contour, then $v_n = \frac{d\sigma}{dl} (\partial_\sigma y \partial_t x - \partial_\sigma x \partial_t y)$, where $dl = |dz| = \sqrt{(dx)^2 + (dy)^2}$ is the line element along the contour and v_n is positive when the contour moves to the right of the increasing σ direction). The Cauchy-type integral in the r.h.s. defines analytic functions both inside and outside the contour. For z inside the contour, choosing a neighborhood of 0 such that $|z| < |z'|$ for all $z' \in \partial \mathbb{D}$, we can expand $M_+(z)$ as in (3.4) and find that

$$\partial_t M_+(z) = \sum_{k=1}^{\infty} k \partial_t T_k z^{k-1} = 0$$

for all z in this neighborhood. By uniqueness of analytic continuation $\partial_t M_+(z) = 0$ everywhere in \mathbb{D} . According to the property of integrals of Cauchy type this means that the function $\partial_t M_+(z)$ is analytic in $\mathbb{C} \setminus \mathbb{D}$ and is given there by the Cauchy integral (3.11). The boundary value of this function is $\text{sign}(y) v_n(z) |dz|/dz$ with *real* $v_n(z)$ almost everywhere on the contour (in our case actually everywhere except maybe the two corner points on the real axis). Furthermore, deformations of symmetric domains preserving symmetry with respect to the real axis obey the condition

$$\oint_{\partial \mathbb{D}} \text{sign}(y) v_n(z) |dz| = 0$$

(because $v_n(z) = v_n(\bar{z})$), which means, according to the Cauchy integral representation, that the function $\partial_t M_+(z)$ has a zero at infinity of at least second order. Invoking the technique from the theory of boundary values of analytic functions [16, 17], one can prove (see Appendix B) that any analytic function with such properties in $\mathbb{C} \setminus \mathbb{D}$ must be identically zero. Therefore, $v_n \equiv 0$, i.e., the deformation is trivial.

3.4 Special deformations of fat slits

In order to define deformations that change one of the moments T_k keeping all other fixed we need the odd Green's function for symmetric domains introduced in Section 2.2.

Fix a point $a \in \mathbb{H} \setminus \mathbb{B}$ and consider small deformations δ_a^- of a fat slit \mathbb{B} defined by the infinitesimal normal displacement of the boundary as follows:

$$\delta_a^- n(\xi) = -\frac{\epsilon}{2} \partial_n G^-(a, \xi), \quad \xi \in \gamma, \quad a \in \mathbb{H} \setminus \mathbb{B}. \quad (3.12)$$

(here $\xi \in \partial\mathbb{D}$ and $\epsilon \rightarrow 0$). These are analogs of the deformations $\delta_a n(\xi) = -\frac{\epsilon}{2} \partial_n G(a, \xi)$ from [14] which generate dToda flows in the space of compact domains in the plane. As we shall soon see, δ_a^- generate, in the same sense, dKP-flows in the space of fat slits. Extending the definition of the δ_a^- to the lower half plane, we can define the corresponding deformation of the symmetric domain $\mathbb{D} = \mathbb{B} \cup \bar{\mathbb{B}}$:

$$\delta_a^- n(\xi) = -\frac{\epsilon}{2} \text{sign}(\text{Im}\xi) \partial_n G^-(a, \xi), \quad \xi \in \partial\mathbb{D}. \quad (3.13)$$

Clearly, $\delta_a^- n(\xi)/\epsilon$ as $\epsilon \rightarrow 0$ is to be understood as normal velocity of the boundary under the deformation.

An important comment is in order. For the deformations δ_a^- to be well defined around the points x_-, x_+ , we assume that the angles α_\pm between the curve γ and the real axis are strictly acute: $0 < \alpha_\pm < \pi/2$ (see Fig. 2). Since $p'(z) \sim (z - x_\pm)^{\frac{\pi}{\alpha_\pm} - 1}$ around the corner points (for a rigorous proof see, e.g., [18, Lemma 2.8]), it is seen from (2.8) that the normal velocity of the boundary near the corner points tends to zero as $\xi \rightarrow x_\pm$ and, moreover, so does the angular velocity of the parts of the boundary near the corners (which is of order $|p''(z)|$). This means that the points x_\pm and the angles α_\pm remain fixed. For not strictly acute angles α_+, α_- the situation is much more complicated. For example, the angles can immediately jump to other values and the deformations are not always well defined (cf. [19]). This case deserves further investigation.

Expanding the Green's function as in (2.10), one can introduce the deformations

$$\delta^{(k,-)} n(\xi) = \frac{\epsilon}{2} \text{sign}(\text{Im}\xi) \partial_n \text{Im} B_k(p(\xi)), \quad \xi \in \partial\mathbb{D}. \quad (3.14)$$

Like δ_a^- , the deformations $\delta^{(k,-)}$ do not shift the endpoints of γ .

It is not difficult to show that $\delta^{(k,-)}$ changes the harmonic moment T_k keeping all other fixed. Indeed, assuming that $a \in \mathbb{D}$, we write:

$$\delta^{(k,-)} M_+(a) = \frac{1}{\pi i} \oint_{\partial\mathbb{D}} \frac{\text{sign}(y) \delta^{(k,-)} n(z)}{a - z} |dz| = \frac{\epsilon}{2\pi i} \oint_{\partial\mathbb{D}} \frac{\partial_n \text{Im} B_k(p(z))}{a - z} |dz|,$$

where the extra sign in the definition of $\delta^{(k,-)}$ cancels the $\text{sign}(y)$ in the definition of $M_+(a)$. Because $\text{Im} B_k(p(z)) = 0$ on $\partial\mathbb{D}$, we have:

$$\delta^{(k,-)} M_+(a) = \frac{\epsilon}{2\pi i} \oint_{\partial\mathbb{D}} \frac{dB_k(p(z))}{z - a} = \frac{\epsilon}{2\pi i} \oint_{\partial\mathbb{D}} \frac{dz^k}{z - a} = \epsilon k a^{k-1},$$

where we have used the fact that the function $B_k - z^k$ is analytic in $\mathbb{C} \setminus \mathbb{D}$ and vanishes at ∞ , and so does not contribute to the integral. We thus see that $\delta^{(k,-)} T_j = \epsilon \delta_{kj}$.

3.5 Vector fields in the space of fat slits

Deformations which depend on γ in a smooth way can be represented by vector fields in the space of fat slits. Let $\delta n(z)$ be any small deformation of a fat slit. Given a functional X on the space of fat slits, its variation reads:

$$\delta X = \int_{\gamma} \frac{\delta X}{\delta n(\xi)} \delta n(\xi) |d\xi|.$$

The variational derivative $\delta X / \delta n(\xi)$ has the following meaning:

$$\frac{\delta X}{\delta n(\xi)} = \lim_{\varepsilon \rightarrow 0} \frac{\delta^{(\varepsilon)} X}{\varepsilon}.$$

Here $\delta^{(\varepsilon)} X$ is the variation of the functional under attaching a small bump of area $\varepsilon \rightarrow 0$ at the point $\xi \in \gamma$ (a symmetric bump is assumed to be attached at the point $\bar{\xi}$). Let $\delta n(\xi) = \varepsilon g(\xi)$, $\varepsilon \rightarrow 0$, then we define the vector field (the Lie derivative) $\nabla^{(g)}$:

$$\nabla^{(g)} X = \int_{\gamma} \frac{\delta X}{\delta n(\xi)} g(\xi) |d\xi|.$$

Applying these general formulas to the deformations δ_a^- (see (3.12)), we can write $\delta_a^- X = \varepsilon \nabla^-(a) X$, where the vector field $\nabla^-(a)$ acts on functionals as follows:

$$\nabla^-(a) X = -\frac{1}{2} \int_{\gamma} \frac{\delta X}{\delta n(\xi)} \partial_n G^-(a, \xi) |d\xi|. \quad (3.15)$$

This equation gives an invariant definition of the vector field $\nabla^-(a)$ independent of any choice of coordinates. According to the Dirichlet formula (2.5), the action of $\nabla^-(a)$ provides the harmonic extension of the function $\pi \delta X / \delta n(\xi)$ from γ to \mathbb{H} bounded at infinity and equal to 0 on the rays of the real axis outside \mathbf{B} .

In the local coordinates T_k , $\nabla^-(a)$ is represented as an infinite linear combination of the vector fields $\partial / \partial T_k$ which can be thought of as partial derivatives. To find it explicitly, we calculate

$$\delta_a^- T_k = -\frac{2}{\pi k} \int_{\gamma} \operatorname{Im}(z^{-k}) \delta_a^- n(z) |dz| = \frac{\varepsilon}{\pi k} \int_{\gamma} \operatorname{Im}(z^{-k}) \partial_n G^-(a, z) |dz| = -\frac{2\varepsilon}{k} \operatorname{Im}(a^{-k}),$$

where the last equality follows from the Dirichlet formula (2.5) for symmetric domains. Now, given a functional X on the space of fat slits, and assuming that X is a function of the moments T_k only, we write

$$\delta_a^- X = \sum_{k \geq 1} \frac{\partial X}{\partial T_k} \delta_a^- T_k = -2\varepsilon \sum_{k \geq 1} \frac{1}{k} \operatorname{Im}(a^{-k}) \frac{\partial X}{\partial T_k} = \varepsilon \nabla^-(a) X,$$

so $\nabla^-(a)$ is given by

$$\nabla^-(a) = -2 \sum_{k \geq 1} \frac{1}{k} \operatorname{Im}(a^{-k}) \frac{\partial}{\partial T_k}. \quad (3.16)$$

The vector fields $\nabla^-(a)$ are “half-plane” analogs of the vector fields $\nabla(a)$ introduced in [14] via their action on functionals X in the space of all domains:

$$\nabla(a) X = -\frac{1}{2} \int_{\partial \mathbf{D}} \frac{\delta X}{\delta n(\xi)} \partial_n G(a, \xi) |d\xi| \quad (3.17)$$

(cf. (3.15)). Below we will show that the dKP hierarchy is related to the vector fields $\nabla^-(a)$ in the same way as the dToda hierarchy is related to the $\nabla(a)$.

4 The dispersionless KP hierarchy

4.1 Lax equations

The dispersionless KP (dKP) hierarchy is encoded in the Hadamard formula (2.6). To see this, fix three points $a, b, c \in \mathbb{H} \setminus \mathbb{B}$ and find $\delta_c^- G^-(a, b)$:

$$\delta_c^- G^-(a, b) = -\frac{\epsilon}{4\pi} \int_{\gamma} \partial_n G^-(a, z) \partial_n G^-(b, z) \partial_n G^-(c, z) |dz|. \quad (4.1)$$

This formula shows that the quantity $\nabla^-(a)G^-(b, c)$ is *symmetric* with respect to all three arguments:

$$\nabla^-(a)G^-(b, c) = \nabla^-(b)G^-(a, c) = \nabla^-(c)G^-(a, b). \quad (4.2)$$

Using the expansions (2.10) and (3.16), we get

$$\frac{\partial}{\partial T_k} \mathcal{I}m(B_l(p(z))) = \frac{\partial}{\partial T_l} \mathcal{I}m(B_k(p(z))), \quad k, l \geq 1.$$

Since $\overline{B_k(p(z))} = B_k(p(\bar{z}))$, we can easily separate holomorphic and antiholomorphic parts of this equality and rewrite it as a relation between functions of z only:

$$\frac{\partial B_l(p(z))}{\partial T_k} = \frac{\partial B_k(p(z))}{\partial T_l}. \quad (4.3)$$

In particular,

$$\frac{\partial B_k(p(z))}{\partial T_1} = \frac{\partial p(z)}{\partial T_k}. \quad (4.4)$$

Treating p rather than z as an independent variable and passing to the inverse map, $z(p)$, one can bring this equality to the form

$$\frac{\partial z(p)}{\partial T_k} = \frac{\partial B_k(p)}{\partial p} \frac{\partial z(p)}{\partial T_1} - \frac{\partial B_k(p)}{\partial T_1} \frac{\partial z(p)}{\partial p} \equiv \{B_k(p), z(p)\}. \quad (4.5)$$

Recalling that $B_k(p) = (z^k(p))_{\geq 0}$ (see (2.12)), we recognize the standard Lax equations of the dKP hierarchy.

So far we assumed that p does not belong to the segment $[p_-, p_+]$ (see Fig. 3). This segment is a branch cut of the function $z(p)$. On this cut we can write

$$z(p \pm i0) = x(p) \pm i|y(p)|, \quad p_- < p < p_+. \quad (4.6)$$

Here $|y(p)| = y(p + i0) = -y(p - i0) \geq 0$. Outside the cut, the real-valued function $x(p)$ has the same expansion (2.2) as the $z(p)$. On the cut, it is given by the principal value integral

$$x(p) = p + \frac{1}{\pi} \text{P.V.} \int_{p_-}^{p_+} \frac{|y(p')| dp'}{p' - p}. \quad (4.7)$$

Because $B_k(p)$ is a polynomial with real coefficients, one sees from (4.5) that $x(p)$ and $|y(p)|$ obey the same Lax equations:

$$\frac{\partial x(p)}{\partial T_k} = \{B_k(p), x(p)\}, \quad \frac{\partial |y(p)|}{\partial T_k} = \{B_k(p), |y(p)|\}, \quad p \in [p_-, p_+]. \quad (4.8)$$

In the next subsection we show that $2|y(p)|$ is the Orlov-Shulman function.

4.2 The Orlov-Shulman function

Consider the functions $M_+(z)$, $M_-(z)$ defined by the integrals of Cauchy type (3.4), (3.10) for z inside and outside the domain $D = B \cup \bar{B}$ respectively. They can be also represented by the Taylor series

$$M_+(z) = \sum_{k \geq 1} k T_k z^{k-1}, \quad M_-(z) = \sum_{k \geq 1} V_k z^{-k-1}$$

which converge in some neighborhoods of 0 and ∞ respectively. The function $M_+(z)$ is analytic everywhere in D while $M_-(z)$ is analytic everywhere in $\mathbb{C} \setminus D$ (with zero of second order at ∞). Moreover, for analytic arcs γ both $M_+(z)$ and $M_-(z)$ can be analytically continued across the arcs γ and $\bar{\gamma}$ everywhere except their endpoints on the real axis, where both functions have a singularity. Therefore, the function

$$M(z) := M_+(z) - M_-(z) = \sum_{k \geq 1} k T_k z^{k-1} - \sum_{k \geq 1} V_k z^{-k-1} \quad (4.9)$$

is analytic in a neighborhood of the boundary of D (excluding the points x_{\pm} on the real axis) and, by the property of the Cauchy-type integrals, is equal to $2|\mathcal{I}m z| = 2|y|$ on $\gamma \cup \bar{\gamma}$. This can be also seen from formulas (3.6), (3.8) (together with their extensions to the lower half plane) taking into account that the derivatives of the electrostatic potential are continuous at the boundary:

$$\partial_z \Phi^-(z) \Big|_{\text{in}} = 2iy \mp iM_+(z) = \mp iM_-(z) = \partial_z \Phi^-(z) \Big|_{\text{out}}$$

where the upper (lower) sign is taken for z in the upper (lower) half plane. We see that for z in the upper half plane $M(z)$ is the analytic continuation of the function $2y$ from the contour γ , while for z in the lower half plane $M(z)$ is the analytic continuation of the function $-2y$ from the complex conjugate contour $\bar{\gamma}$, i.e., $M(z(p)) = 2|y(p)|$.

Let $S(z)$ be the Schwarz function of the contour γ , i.e., an analytic function such that $S(z) = \bar{z}$ for z on γ (see [20] for details). For analytic contours, it is known to be well defined in some strip-like neighborhood of the curve. Clearly, the Schwarz of the complex conjugate contour $\bar{\gamma}$ is then $\bar{S}(z) = \overline{S(\bar{z})}$. By uniqueness of analytic continuation, we can express $M(z)$ in terms of the Schwarz function:

$$M(z) = \begin{cases} i(S(z) - z), & \text{Im } z > 0 \\ -i(\bar{S}(z) - z), & \text{Im } z < 0. \end{cases} \quad (4.10)$$

Let us show that

$$\partial_{T_k} M(z) = \partial_z B_k(p(z)). \quad (4.11)$$

Consider the change of the $S(z)$ under the deformation δ_a^- . If $z \in \gamma$, then, using the identity $v_n(z) = \frac{\partial_T S(z)}{2i\sqrt{S'(z)}}$ for the normal velocity of the boundary under a deformation

with a parameter T , we can write

$$\delta_a^- S(z) = -2i\sqrt{S'(z)} \frac{\epsilon}{2} \partial_n G^-(a, z).$$

Since $\sqrt{S'(z)} = |dz|/dz = 1/\tau(z)$, where $\tau(z)$ is the unit tangent vector to the curve γ (represented as a complex number) and $\partial_n G^-(a, z) = -2i\tau(z)\partial_z G^-(a, z)$, we have

$$\delta_a^- S(z) = -2\epsilon\partial_z G^-(a, z)$$

and so,

$$\nabla^-(a)S(z) = -2\partial_z G^-(a, z) \quad (4.12)$$

for $z \in \gamma$ and, by analytic continuation, everywhere in the neighborhood where $S(z)$ is a well defined analytic function. Expanding both sides as in (2.10), (3.16), we finally get

$$\partial_{T_k} S(z) = -i\partial_z B_k(p(z)), \quad (4.13)$$

which is equivalent to (4.11).

An important particular case of (4.11) is

$$\partial_{T_1} M(z) = \partial_z p(z). \quad (4.14)$$

Passing to partial derivatives at constant p , one can rewrite it in the form of the “string equation”:

$$\{z(p), 2|y(p)|\} = 1, \quad p \in [p_-, p_+]. \quad (4.15)$$

This relation together with the Lax equations (4.8) show that $M(z(p)) = 2|y(p)|$ is the Orlov-Shulman function [21] of the dKP hierarchy which describes deformations of fat slits.

A closely related useful object is the indefinite integral of $M(z)$. Using the notation of the previous subsection, we introduce the function

$$\begin{aligned} \Omega(z) &= \int_0^z M_+(z)dz + \int_z^\infty M_-(z)dz \\ &= \sum_{k \geq 1} T_k z^k + \sum_{k \geq 1} \frac{V_k}{k} z^{-k}. \end{aligned} \quad (4.16)$$

It is analytic in the same strip-like neighborhood of the curve γ where the Schwarz function is well defined. Since the electrostatic potential $\Phi^-(z)$ is continuous on γ , it follows from (3.7), (3.8) that

$$\text{Im} \Omega(z) = y^2, \quad z \in \gamma. \quad (4.17)$$

The real part of $\Omega(z)$ at $z \in \gamma$ has the meaning of the partial area beneath the curve γ . More precisely, let

$$A(z) = \int_{x_-}^x y' dx', \quad z = x + iy \in \gamma$$

be the partial area of \mathbf{B} cut from the right by a line orthogonal to the real axis and passing through z , then

$$\text{Re} \Omega(z_1) - \text{Re} \Omega(z_2) = 2(A(z_1) - A(z_2)), \quad z_1, z_2 \in \gamma. \quad (4.18)$$

By construction, partial T_k -derivatives of the function Ω at constant z are the generators of the flows:

$$B_k(p(z)) = \partial_{T_k} \Omega(z). \quad (4.19)$$

In this sense the function $\Omega = \Omega(z; \{T_j\})$ solves the whole set of equations (4.3) and thus provides a solution to the dKP hierarchy.

4.3 The string equation

The string equation can be also derived in a different way along the lines of [13]. The idea is to use equation (4.2) with one of the three points lying on the contour γ :

$$\nabla^-(\xi)G^-(a, b) = \nabla^-(a)G^-(b, \xi), \quad \xi \in \gamma,$$

and the other two points tending to infinity. From (3.15) we see that the l.h.s. is equal to $\frac{1}{2}\partial_n G^-(a, \xi)\partial_n G^-(b, \xi)$ which is $2\mathcal{I}m(a^{-1})\mathcal{I}m(b^{-1})(\partial_n \mathcal{I}m p(\xi))^2$ as $a, b \rightarrow \infty$. The r.h.s. in the same limit is $-4\mathcal{I}m(a^{-1})\mathcal{I}m(b^{-1})\partial_{T_1} \mathcal{I}m p(\xi)$. Equating them, we obtain the important relation

$$2\partial_{T_1} \mathcal{I}m p(z) = -|\partial_z p(z)|^2, \quad z \in \gamma. \quad (4.20)$$

Its extension to the lower half of the boundary of D reads

$$2\partial_{T_1} \mathcal{I}m p(z) = |\partial_z p(z)|^2, \quad z \in \bar{\gamma}. \quad (4.21)$$

Passing to the variable p with the help of the identity

$$\partial_{T_1} z(p) = -\frac{\partial_{T_1} p(z)}{\partial_z p(z)}$$

we rewrite equations (4.20), (4.21) in the form

$$\frac{\partial z(p - i\epsilon)}{\partial p} \frac{\partial z(p + i\epsilon)}{\partial T_1} - \frac{\partial z(p - i\epsilon)}{\partial T_1} \frac{\partial z(p + i\epsilon)}{\partial p} = i \operatorname{sign} \epsilon, \quad \epsilon \rightarrow 0. \quad (4.22)$$

Note that the equation at $\epsilon < 0$ is obtained from the one at $\epsilon > 0$ by complex conjugation. Passing to the functions $x(p)$ and $y(p)$, we see that (4.22) coincides with (4.15).

Eq. (4.22) can be cast into the form of an evolution equation for $z(p)$. To derive it, we start from the Hadamard formula for the deformation $\delta^{(1,-)}$:

$$\delta^{(1,-)} G^-(a, z) = \frac{\epsilon}{4\pi} \int_{\gamma} \partial_n G^-(a, \xi) \partial_n G^-(z, \xi) \partial_n \mathcal{I}m p(\xi) |d\xi|.$$

Expanding the Green's function at $|a| \rightarrow \infty$ (see (2.10)) and using the fact that $\delta^{(1,-)} T_1 = \epsilon$, $\delta^{(1,-)} T_k = 0$ at $k \geq 2$, we rewrite it as

$$\partial_{T_1} \mathcal{I}m p(z) = \frac{1}{4\pi} \int_{\gamma} \partial_n G^-(z, \xi) |p'(\xi)|^2 |d\xi|,$$

which, after extracting the holomorphic part and passing to the integration in the p -plane, reads

$$\partial_{T_1} p(z) = \frac{1}{2\pi} \int_{p_-}^{p_+} \frac{dp}{(p(z) - p)|z'(p)|^2}.$$

In terms of the function $z(p)$ we have:

$$\partial_{T_1} z(p) = \frac{z'(p)}{2\pi} \int_{p_-}^{p_+} \frac{dp'}{(p' - p)|z'(p')|^2}, \quad (4.23)$$

which is the desired evolution equation equivalent to (4.22). The latter is obtained from (4.23) by taking the jump of both sides across the segment $[p_-, p_+]$. Because the function $z(p)$ is analytic in the upper half plane, this is equivalent to the full equation (4.23).

5 The dKP hierarchy in the Hirota form

Here we reformulate the dKP hierarchy in the Hirota form using the dispersionless “tau-function” (free energy) and clarify the geometric meaning of the latter.

5.1 Functional F^- and its variations

Given a domain D , not necessarily symmetric, one can introduce the functional F :

$$F = -\frac{1}{\pi^2} \int_D \int_D \log |z^{-1} - \zeta^{-1}| d^2 z d^2 \zeta. \quad (5.1)$$

If the boundary is analytic, F is the “tau-function for analytic curves” introduced in [12] (more precisely, a properly rescaled logarithm of the dToda tau-function). In the 2D electrostatic interpretation, it has the meaning of the electrostatic energy of a uniformly charged domain D with a compensating point-like charge at the origin. It was found in [12] that the Green’s function $G(a, b)$ is given by

$$G(a, b) = \log |a^{-1} - b^{-1}| + \frac{1}{2} \nabla(a) \nabla(b) F, \quad (5.2)$$

where the vector field $\nabla(z)$ in the space of all domains is defined in (3.17) (see [13, 14] for more details). We are going to derive an analog of equation (5.2) for fat slits in the upper half plane.

Given a fat slit B , consider the following functional:

$$F^- = -\frac{1}{\pi^2} \int_B \int_B \log \left| \frac{z - \zeta}{z - \bar{\zeta}} \right| d^2 z d^2 \zeta. \quad (5.3)$$

It has the meaning of the 2D electrostatic energy of the uniformly charged fat slit in the presence of a conductor placed along the real axis. Taking a variation of F^- , it is easy to find how the vector field $\nabla^-(a)$ acts on F^- . We use the general relations given in section 4.2. We have:

$$\pi \frac{\delta F^-}{\delta n(\xi)} = -\frac{2}{\pi} \int_B \log \left| \frac{z - \xi}{z - \bar{\xi}} \right| d^2 z.$$

The function in the r.h.s. is already harmonic and bounded in $\mathbb{H} \setminus B$ as it stands, hence

$$\nabla^-(a) F^- = -\frac{2}{\pi} \int_B \log \left| \frac{a - z}{a - \bar{z}} \right| d^2 z = \Phi^-(a). \quad (5.4)$$

Expanding both sides as $a \rightarrow \infty$ and comparing coefficients in front of the basis harmonic functions $\mathcal{I}m(a^{-k})$, we find:

$$\frac{\partial F^-}{\partial T_k} = \frac{2}{\pi} \mathcal{I}m \int_B z^k d^2 z = V_k, \quad (5.5)$$

where V_k are the interior harmonic moments. Proceeding in a similar way, we find

$$\pi \frac{\delta \Phi^-(a)}{\delta n(\xi)} = -2 \log \left| \frac{\xi - a}{\xi - \bar{a}} \right|.$$

The r.h.s. is harmonic (in ξ) everywhere in $\mathbb{H} \setminus \mathbb{B}$ except the point a . This singularity can be canceled by adding the Green's function G^- (which vanishes on the boundary). Therefore,

$$\nabla^-(b)\Phi^-(a) = -2 \log \left| \frac{a-b}{a-\bar{b}} \right| + 2G^-(a, b),$$

and we obtain the formula for $G^-(a, b)$,

$$G^-(a, b) = \log \left| \frac{a-b}{a-\bar{b}} \right| + \frac{1}{2} \nabla^-(a) \nabla^-(b) F^-, \quad (5.6)$$

which is a “half-plane” analog of (5.2).

5.2 Hirota equations for the dKP hierarchy

Equation (5.2) is known to encode the dToda hierarchy in the Hirota form. Equation (5.6) does the same for the dKP hierarchy. To see this, let us apply the arguments from [14].

Combining (5.6) and (2.7), we obtain the relation

$$\log \left| \frac{p(z) - p(z')}{p(z) - p(\bar{z}')} \right|^2 = \log \left| \frac{z - z'}{z - \bar{z}'} \right|^2 + \nabla^-(z) \nabla^-(z') F^-, \quad (5.7)$$

which implies an infinite hierarchy of differential equations for the function F^- . Recall that the conformal map $p(z)$ is normalized as

$$p(z) = z + \frac{u}{z} + O(1/z^2), \quad z \rightarrow \infty \quad (5.8)$$

(see (2.1)). Tending $z' \rightarrow \infty$ in (5.7), one gets:

$$\mathcal{I}m p(z) = \mathcal{I}m z + \frac{1}{2} \partial_{T_1} \nabla^-(z) F^-. \quad (5.9)$$

The limit $z \rightarrow \infty$ of this equality yields a simple formula for the capacity:

$$u = \partial_{T_1}^2 F^-. \quad (5.10)$$

Let us separate holomorphic parts of these equations, introducing the holomorphic part of the operator $\nabla^-(z)$:

$$D(z) = \sum_{k \geq 1} \frac{z^{-k}}{k} \partial_{T_k}, \quad \nabla^-(z) = i [D(z) - D(\bar{z})]. \quad (5.11)$$

Equation (5.7) then implies the relation

$$\log \frac{p(z) - p(z')}{p(z) - p(\bar{z}')} = \log \frac{z - z'}{z - \bar{z}'} + i D(z) \nabla^-(z') F^-, \quad (5.12)$$

which is holomorphic in z . In the limit $z' \rightarrow \infty$ it gives the formula for the conformal map $p(z)$:

$$p(z) = z + \partial_{T_1} D(z) F^- \quad (5.13)$$

(this formula also follows from (5.9)). In a similar way, equation (5.12) implies the relation

$$\log \frac{p(z) - p(z')}{z - z'} = -D(z)D(z')F^-, \quad (5.14)$$

which is holomorphic in both z and z' . Taking into account (5.13), we rewrite it as follows:

$$1 - e^{-D(a)D(b)F^-} = -\frac{D(a) - D(b)}{a - b} \partial_{T_1} F^-. \quad (5.15)$$

It is the dKP hierarchy in the Hirota form. We see that the function F^- is the dispersionless tau-function for this hierarchy. The double integral representation (5.3) clarifies its geometric meaning.

6 A growth model associated with dKP hierarchy

The special deformations from section 4.1 suggest to introduce a growth model which is associated with the dKP hierarchy in the same way as the Laplacian growth [8, 9] of compact planar domains at zero surface tension is associated [5] with the dToda hierarchy. In fact the model to be introduced is also of the Laplacian type, i.e., the interface dynamics is governed by the Darcy law, but differs from the standard one by boundary conditions.

The idea should be already clear from section 4.1: to consider growth of a fat slit under the deformation $\delta^{(1,-)}$ which changes only the first harmonic moment T_1 keeping all other fixed and to identify T_1 with time T .

The corresponding growth problem can be formulated as follows. Consider a fat slit $B(T)$ with a moving boundary $\gamma(T)$, where T is time, and suppose that the motion of the boundary follows the Darcy law:

$$v_n(\xi) = \frac{1}{2} \partial_n \phi(\xi), \quad \xi \in \gamma. \quad (6.1)$$

Here $v_n(\xi) = \delta n(\xi)/\delta T$ is the normal velocity of the boundary at the point ξ and $\phi(z)$ is a harmonic function in $\mathbb{H} \setminus B$ such that

- (i) $\phi = 0$ on γ and on the rays of the real axis $[-\infty, x_-]$, $[x_+, +\infty]$;
- (ii) $\phi(z) = \mathcal{I}m z + o(1)$ as $\mathcal{I}m z \rightarrow +\infty$.

Clearly, $\phi(z) = \mathcal{I}m p(z)$, where $p(z)$ is the conformal map (2.1), is harmonic in $\mathbb{H} \setminus B$ and obeys these conditions. Comparing with (3.14) at $k = 1$, we see that the dynamics is given by the deformation $\delta^{(1,-)}$ at any point in time $T = T_1$ and all the higher moments T_k are integrals of motion. In other words, for our growth process $\partial_T M_+(z) = 1$. Equivalently, the dynamics can be reformulated in terms of the inverse conformal map $z(p, T)$ as the “string equation” in the form (4.22),

$$\mathcal{I}m \left[\partial_p z(p - i0, T) \partial_T z(p + i0, T) \right] = \frac{1}{2}, \quad p \in [p_-, p_+], \quad (6.2)$$




Figure 5: a): The Laplacian growth of a fat slit associated with dKP hierarchy; b) The standard Laplacian growth in the upper half plane.




Figure 6: The curve $|z^2 + T^2| = T^2$ in the upper half plane. The tangent lines at the origin are at an angle of 45° to the real axis.

or in the form of the evolution equation (4.23) (a similar equation for the Laplacian growth in the standard setting is well known [22]). As was already mentioned, the growth process is well defined if both angles α_\pm between γ and the real axis are acute. Then these angles and the points x_- , x_+ stay fixed all the time.

Comparing this setting with the standard Laplacian growth in the upper half plane, we see that the conditions on ϕ are very similar if not the same: $\phi = 0$ on an infinite contour from left to right infinity, harmonic above it and tends to $\mathcal{I}m z$ as $\mathcal{I}m z \rightarrow +\infty$. However, in our case, unlike in the standard one, only a finite part of the boundary (namely, the part which lies above the real axis) moves according to the Darcy law while the remaining part (the rays of the real axis) is kept fixed despite the fact that the gradient of ϕ is nonzero there (Fig. 5). We do not know a proper hydrodynamic realization of this growth process.

So far we assumed that x_- was strictly less than x_+ , so that the base of a fat slit was a segment of nonzero length. The setting of this section allows us to consider the degenerate case $x_- = x_+ = 0$ when the base of a fat slit consists of one point. The first harmonic moment $T_1 = T$ as well as the function $M_+(z)$ are still well-defined but $M_+(z)$ is singular at $z = 0$ and thus can not be expanded into the Taylor series around this point (this means that the higher harmonic moments (3.1) are ill-defined).

The simplest explicit solution to equation (6.2) known to us describes self-similar

growth of a “fat slit” with degenerate base. The function

$$p(z, T) = \frac{z^2 + 2T^2}{\sqrt{z^2 + T^2}} \quad (6.3)$$

performs a conformal map from the exterior in \mathbb{H} of the curve $|z^2 + T^2| = T^2$, or, in polar coordinates,

$$R(\theta) = T\sqrt{-2\cos 2\theta}, \quad \frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}, \quad (6.4)$$

to the upper half plane. This curve is shown in Fig. 6. In this case $x_- = x_+ = 0$, $\alpha_- = \alpha_+ = \pi/4$, $p_{\pm} = \pm 2T$. The inverse map has the form

$$z(p, T) = \frac{1}{\sqrt{2}}(p^2 - 4T^2)^{1/4} \left(p + (p^2 - 4T^2)^{1/2} \right)^{1/2}. \quad (6.5)$$

One can check that it does solve equation (6.2). More results on explicit solutions to Laplacian growth of fat slits will be published elsewhere.

7 Concluding remarks

We have constructed a parametric family of conformal maps of the upper half plane which is related to the dKP hierarchy with real “times” T_k in the same way as conformal maps of the unit disk onto compact domains in the plane with smooth boundary are related to the dToda hierarchy with complex conjugate “times” t_k, \bar{t}_k . Like in the dToda case, the deformations of domains (“fat slits”) in the upper half plane induced by dKP flows have a physical interpretation as Laplacian growth with certain type of sources or sinks at infinity. At the same time, our construction extends the well known connection between the dKP hierarchy and conformal maps of slit domains. In all cases, the conformal map plays the role of the Lax function.

However, a limiting procedure from fat slits to usual slits is singular and is not easy to trace on the level of the Lax equations. We hope that a better understanding of this limit will further clarify the geometric meaning of solutions to equations of the dKP hierarchy. We also expect that yet more general solutions can be obtained by the same method applied to the case of a background charge distributed in the upper half plane with a non-uniform density, in accordance with a similar construction given in [23].

The solutions discussed in this paper have a nice geometric meaning but it seems to be very hard to express them analytically in a closed form. In this respect the situation is less favourable than in the dToda case, where some simple explicit solutions for conformal maps as functions of a finite number of nonzero harmonic moments (corresponding, for example, to a parametric family of ellipses) are available. It is clear that in our case the situation when only a finite number of the moments T_k are nonzero can not be realized because the local behavior of their generating function $M_+(z)$ near the points x_- , x_+ (which can be found from the integral representation (3.4)) shows that they are branch points of this function. This suggests that the corresponding solutions may be similar to the multi-cut solutions to Laplacian growth discussed recently in [24].

8 Appendices

Appendix A

Here we give some details of the derivation of the formula (3.11) for the time derivative of the function $M_+(z)$:

$$\partial_t M_+(a) = \frac{1}{\pi i} \oint_{\partial D} \frac{\text{sign}(y)v_n(z)}{a-z} |dz|. \quad (8.1)$$

Here $v_n(z)$ is the velocity of the normal displacement of the boundary at the point z . Let $z(\sigma, t) = x(\sigma, t) + iy(\sigma, t)$ be any parametrization of the contour such that σ is a steadily increasing function of the arc length, then it is a simple kinematical fact that

$$v_n = \frac{d\sigma}{dl} (\partial_\sigma y \partial_t x - \partial_\sigma x \partial_t y), \quad (8.2)$$

where $dl = |dz| = \sqrt{(dx)^2 + (dy)^2}$ is the line element along the contour. According to our convention, $v_n(z)$ is positive when the contour, in a neighborhood of the point z , moves to the right of the increasing σ direction.

Without loss of generality, we assume that σ varies from 0 to 2π , and, furthermore, for symmetric contours we choose σ in such a way that $z(2\pi - \sigma, t) = \overline{z(\sigma, t)}$, $z(0, t) = z(2\pi, t) = x_+$, $z(\pi, t) = x_-$ and $y(\sigma, t)$ is positive for $0 < \sigma < \pi$. We have: $M_+(a) = I(a) + \overline{I(\bar{a})}$, where

$$I(a) = \frac{1}{\pi i} \int_0^\pi \frac{y(\sigma) dz(\sigma)}{z(\sigma) - a}.$$

A straightforward calculation gives:

$$\begin{aligned} i\pi \partial_t I(a) &= \int_0^\pi \left(\frac{\partial_t y \partial_\sigma z + y \partial_t \partial_\sigma z}{z - a} - \frac{y \partial_t z \partial_\sigma z}{(z - a)^2} \right) d\sigma \\ &= \int_0^\pi \left(\frac{\partial_t y \partial_\sigma z + y \partial_t \partial_\sigma z}{z - a} d\sigma + y \partial_t z d\left(\frac{1}{z - a}\right) \right) \\ &= \int_0^\pi \frac{\partial_t y \partial_\sigma z - \partial_t z \partial_\sigma y}{z - a} d\sigma + \frac{y(\sigma) \partial_t z(\sigma)}{z(\sigma) - a} \Big|_0^\pi. \end{aligned}$$

The last term obviously vanishes and we obtain, using (8.2):

$$\partial_t I(a) = \frac{1}{\pi i} \int_\gamma \frac{v_n(z) |dz|}{a - z}.$$

Adding $\overline{\partial_t I(\bar{a})}$, we get (8.1).

Appendix B

In this Appendix we outline the proof of the following proposition.

Proposition 1. Let D be a compact domain bounded by a closed piecewise analytic contour $\Gamma = \partial D$ in the plane with a finite number of corner points. Consider the function $h(z)$ defined by the Cauchy-type integral

$$h(z) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{\rho(\xi) |d\xi|}{z - \xi}, \quad (8.3)$$

where $\rho(\xi)$ is a bounded real-valued piecewise continuous function on Γ such that

$$\oint_{\Gamma} \rho(\xi) |d\xi| = 0 \quad (8.4)$$

and assume that $h(z) = 0$ for all $z \in D$. Then $\rho \equiv 0$.

One can try to prove this statement by means of the following elementary argument. Let $\tau(\xi) = d\xi/|d\xi|$ be the unit tangential vector to the curve Γ at the point ξ represented as a complex number. If $h(z) = 0$ for all $z \in D$, then the properties of Cauchy-type integrals imply that $\rho(\xi)/\tau(\xi)$ is the boundary value of a holomorphic function $h(z)$ in $\mathbb{C} \setminus D$ vanishing at infinity. In fact $h(z)$ is given by the same integral (8.3), where $z \in \mathbb{C} \setminus D$. Condition (8.4) tells us that the zero at infinity is of at least second order. Let $w(z)$ be the conformal map from $\mathbb{C} \setminus D$ onto the unit disk such that $w(\infty) = 0$ and $r = \lim_{z \rightarrow \infty} zw(z)$ is real positive. By the well known property of conformal maps we have

$$\frac{dz}{|dz|} e^{i \arg w'(z)} = \frac{dw}{|dw|}$$

along the curve Γ . Therefore,

$$\tau(z) = i|w'(z)| \frac{w(z)}{w'(z)}, \quad (8.5)$$

and we thus see that

$$\frac{\rho(z) w'(z)}{i|w'(z)|w(z)}$$

is the boundary value of the holomorphic function $h(z)$. Since $w'(z) \neq 0$ in $\mathbb{C} \setminus D$, the function

$$g(z) = h(z) \frac{w(z)}{w'(z)}$$

is holomorphic there with the *purely imaginary* boundary value

$$\frac{\rho(z)}{i|w'(z)|}.$$

Then the real part of this function is harmonic and bounded in $\mathbb{C} \setminus D$ and is equal to 0 on the boundary. By uniqueness of a solution to the Dirichlet boundary value problem, $\operatorname{Re} g(z)$ must be equal to 0 identically. Therefore, $g(z)$ takes purely imaginary values everywhere in $\mathbb{C} \setminus D$ and so is a constant. By virtue of condition (8.4) this constant must be 0 which means that $\rho \equiv 0$.

However, this argument is directly applicable only for purely analytic contours for which all singularities and zeros of the function $w'(z)$ lie strictly inside it. For contours

with corners, the corner points are singular points of the conformal map $w(z)$. Some more work is required to make the above argument rigorous. Below we present another proof of Proposition 1, which makes use of some non-trivial facts about boundary values of analytic functions and actually works in a much more general setting than just a finite number of corner points¹. It takes advantage of translating the proposition to a statement about analytic functions in the unit disk.

Sketch of proof of Proposition 1. If $h(z)$ defined by (8.3) is identically 0 in \mathbb{D} , then it is analytic in $\mathbb{C} \setminus \mathbb{D}$, is $O(1/z^2)$ near infinity (because of (8.4)) and has the boundary value $\rho(z)/\tau(z)$ almost everywhere on Γ . (In our situation “almost everywhere” means everywhere except a finite number of points.) It is known [16, chapter III] that $h(z)$ belongs to the Smirnov class $E^1(\mathbb{C} \setminus \mathbb{D})$.

Let $z = \varphi(w)$ be the conformal map from the unit disk onto $\mathbb{C} \setminus \mathbb{D}$ such that $\varphi(0) = \infty$ and $\varphi(w) = r/w + O(1)$ as $w \rightarrow 0$ with real r . (The function $\varphi(w)$ is inverse to $w(z)$ introduced above.) Set $\tilde{h}(w) = h(\varphi(w))$. Since

$$\tau(\varphi(w)) = \frac{dz}{|dz|} = \frac{d\varphi(w)}{|d\varphi(w)|} = \frac{\varphi'(w)dw}{|\varphi'(w)||dw|} = iw \frac{\varphi'(w)}{|\varphi'(w)|}$$

it follows from the above that $\tilde{h}(w)$ is analytic in the unit disk with the boundary value

$$\tilde{h}(w) = \frac{\rho(\varphi(w))|\varphi'(w)|}{iw \varphi'(w)} \quad (8.6)$$

almost everywhere on the unit circle and has zero of at least second order at $w = 0$. Clearly, the function $\tilde{H}(w) = \tilde{h}(w)\varphi'(w)$ is analytic in the unit disk because the second order pole of $\varphi'(w)$ at $w = 0$ is canceled by the zero of $\tilde{h}(w)$. Furthermore, according to the Keldysh-Lavrentiev theorem [17, chapter 10], $\tilde{H}(w)$ belongs to the Hardy class H^1 . But then the function $H(w) = w\tilde{H}(w)$ belongs to the same Hardy class and takes *purely imaginary boundary values*

$$-i\rho(\varphi(w))|\varphi'(w)|$$

almost everywhere on the unit circle. The characteristic property of functions from the class H^1 is that they can be represented by the Poisson integral of their boundary values with real positive Poisson kernel (see [16, chapter II, §5] or Theorem 3.9 in [17]). This means that $H(w)$ must be purely imaginary everywhere inside the unit disk and hence must be a constant. Since $H(0) = 0$, the constant is 0, so $h(z)$ vanishes identically and $\rho \equiv 0$.

The proof extends word for word to a more general case when ρ is an integrable function with respect to $|dz|$ (not necessarily bounded) and the boundary of \mathbb{D} is a rectifiable Jordan curve. It is crucial that the differential $\rho(z)|dz|$ is real valued on the boundary. If one dropped that assumption, the statement is false. Moreover, functions from the class E^1 can have real boundary values in domains with cusps (see an example in [25]).

¹I thank D.Khavinson who suggested this proof and explained it to me.

Acknowledgments

The author thanks D.Khavinson, I.Krichever, M.Mineev-Weinstein, T.Takebe, D.Vasiliev and P.Wiegmann for discussions and D.Khavinson for reading the manuscript. He is also grateful to organizers of the workshops “Laplacian growth and related topics” (CRM, Montreal, August 2008) and “Geometry and integrability in mathematical physics” (CIRM, Luminy, September 2008), where these results were reported. This work was supported in part by RFBR grant 08-02-00287, by grant for support of scientific schools NSh-3035.2008.2 and by the ANR project GIMP No. ANR-05-BLAN-0029-01.

References

- [1] J. Gibbons and S. Tsarev, *Reductions of the Benney equations*, Phys. Lett. **A211** (1996) 19-24;
J. Gibbons and S. Tsarev, *Conformal maps and reductions of the Benney equations*, Phys. Lett. **A258** (1999) 263-271.
- [2] P. Wiegmann and A. Zabrodin, *Conformal maps and integrable hierarchies*, Commun. Math. Phys. **213** (2000) 523-538.
- [3] I. Krichever, *The τ -function of the universal Whitham hierarchy, matrix models and topological field theories*, Comm. Pure Appl. Math. **47** (1994) 437-475, e-print archive: hep-th/9205110;
I. Krichever, *Funct. Anal Appl.* **22** (1989) 200-213;
I. Krichever, *The dispersionless Lax equations and topological minimal models*, Commun. Math. Phys. **143** (1991) 415-429.
- [4] K. Takasaki and T. Takebe, *Integrable hierarchies and dispersionless limit*, Rev. Math. Phys. **7** (1995) 743-808;
K. Takasaki and T. Takebe, *SDiff(2) Toda equation – hierarchy, tau function and symmetries*, Lett. Math. Phys. **23** (1991) 205-214.
- [5] M. Mineev-Weinstein, P. Wiegmann and A. Zabrodin, *Integrable structure of interface dynamics*, Phys. Rev. Lett. **84** (2000) 5106-5109, e-print archive: nlin.SI/0001007.
- [6] I. Krichever, M. Mineev-Weinstein, P. Wiegmann and A. Zabrodin, *Laplacian growth and Whitham equations of soliton theory*, Physica **D 198** (2004) 1-28, e-print archive: nlin.SI/0311005.
- [7] A. Zabrodin, *Growth processes related to the dispersionless Lax equations*, Physica **D235** (2007) 101-108, e-print archive: math-ph/0609023.
- [8] A comprehensive list of relevant papers published prior to 1998 can be found in: K. A. Gillow and S. D. Howison, *A bibliography of free and moving boundary problems for Hele-Shaw and Stokes flow*, <http://www.maths.ox.ac.uk/~howison/Hele-Shaw/>
- [9] B. Gustafsson, A. Vasil’ev, *Conformal and Potential Analysis in Hele-Shaw Cells*, Birkhäuser Verlag, 2006.

- [10] L. Yu and J. Gibbons, *The initial value problem for reductions of the Benney equations*, Inverse Problems, **16** (2000) 605-618.
- [11] M. Manas, L. Martinez Alonso and E. Medina, *Reductions and hodograph solutions of the dispersionless KP hierarchy*, J. Phys. A.: Math. Gen., **35** (2002) 401-417;
M. Manas, *S-functions, reductions and hodograph solutions of the r-th dispersionless modified KP and Dym hierarchies*, J. Phys. A.: Math. Gen., **37** (2004) 11191-11221.
- [12] I. Kostov, I. Krichever, M. Mineev-Weinstein, P. Wiegmann and A. Zabrodin, *τ -function for analytic curves*, in: Random Matrix Models and Their Applications, Math. Sci. Res. Inst. Publ. vol. 40, Cambridge University Press, pp. 285-299, e-print archive: hep-th/0005259.
- [13] A. Marshakov, P. Wiegmann and A. Zabrodin, *Integrable Structure of the Dirichlet Boundary Problem in Two Dimensions*, Commun. Math. Phys. **227** (2002) 131-153.
- [14] I. Krichever, A. Marshakov and A. Zabrodin, *Integrable Structure of the Dirichlet Boundary Problem in Multiply-Connected Domains*, Commun. Math. Phys. **259** (2005) 1-44.
- [15] J. Hadamard, *Mém. présentés par divers savants à l'Acad. sci.*, **33** (1908).
- [16] I. I. Privalov, *Randeigenschaften Analytischer Funktionen*, Deutscher Verlag Wissenschaften, East Berlin, 1956 (translated from the second Russian edition I. I. Privalov, *Granichnie svoistva analiticheskikh funkciy*, Gostehizdat, Moscow, Leningrad, 1950).
- [17] P. Duren, *Theory of H^p spaces*, Academic Press, New York, 1970.
- [18] J. Akeroyd, D. Khavinson and H. Shapiro, *Remarks concerning cyclic vectors in Hardy and Bergman spaces*, Michigan Math. J. **38** (1991) 191-205.
- [19] J. R. King, A. A. Lacey and J. L. Vazquez, *Persistence of corners in free boundaries in Hele-Shaw flow*, Euro. J. Appl. Math. **6** (1995) 455-490.
- [20] P. J. Davis, *The Schwarz function and its applications*, The Carus Math. Monographs, No. 17, The Math. Assotiation of America, Buffalo, N.Y., 1974.
- [21] A. Orlov and E. Shulman, *Additional symmetries for integrable equations and conformal algebra representation*, Lett. Math. Phys. **12** (1986) 171-179.
- [22] B. Shraiman and D. Bensimon, *Singularities in nonlocal interface dynamics*, Phys. Rev. **A30** (1984) 2840-2842.
- [23] A. Zabrodin, *Dispersionless limit of Hirota equations in some problems of complex analysis*, Theor. and Math. Phys. **129** (2001) 239-257.
- [24] Ar. Abanov, M. Mineev-Weinstein and A. Zabrodin, *Multi-cut solutions to Laplacian growth*, submitted to Physica D.
- [25] D. Khavinson, *Remarks concerning boundary properties of analytic functions of E_p -classes*, Indiana Univ. Math. J. **31** (1982) 779-787.